



Biological invasions: Prospects for slowing a major global change

Daniel Simberloff^{1*}

¹Department of Ecology & Evolutionary Biology, University of Tennessee, Knoxville, Knoxville, Tennessee, United States

*dsimberloff@utk.edu

One of the greatest ongoing ecological changes of the Anthropocene is the rearrangement of the earth's biogeography, as people deliberately or inadvertently move species around. We have been doing this for a long time – the Lapita people from southeast Asia brought various plants and animals (including the Pacific rat) to many islands in the Pacific beginning about 3,000 years ago. However, the number of species introduced to new regions has massively increased in the last few decades as the forces that transport them have grown. From 1970 to 2010, the amount of cargo shipped by sea tripled, and it is on course to double once again by 2030. Amount of air cargo is growing even more rapidly, and passenger air travel is increasing at almost as high a rate. We could as well call the Anthropocene the Homogeocene.

The array of impacts of introduced species is staggering. Most obvious are those of introduced predators on native prey (e.g., rats on many island seabird populations [Pascal, 2011]) and introduced herbivores on native island vegetation (e.g., Cronk, 1989). The spread of the Burmese python (*Python molurus bivittatus*) in Florida and the subsequent drastic decline of populations of prey species ranging in size from rabbits to bobcats (Dorcas et al., 2012) is a noteworthy recent example. Introduced species can compete with native species and also transmit new pathogens to them. The North American gray squirrel (*Sciurus carolinensis*) does both to the European red squirrel (*S. vulgaris*), outcompeting it for nuts and spreading the parapox virus, to which the gray squirrel is resistant and the red squirrel is highly susceptible (Rushton et al., 2006). The red squirrel population has declined drastically in Great Britain, and the gray squirrel is now spreading on the continent from an initial escape from captivity in Italy (Wauters, Tosi and Gurnell, 2002). Competition can be between unrelated species – in New Zealand, the introduced wasp *Vespula vulgaris* outcompetes threatened native birds for “honeydew” produced by scale insects (Beggs and Wardle, 2006). Introduced parasites and pathogens devastate native species. Examples include avian malaria, a major threat to many native Hawaiian birds (van Riper et al., 1986) after its introduction with Asian caged songbirds; crayfish plague, introduced to Europe with resistant North American crayfish and currently ravaging native crayfish (Lodge et al., 2012); and white-nose disease, a European fungus threatening several North American bat species (Turner, Reeder and Coleman, 2011). Some native species undergo a kind of “genetic extinction” from hybridization with related invaders. For instance, North American mallards (*Anas platyrhynchos*) introduced for hunting have extensively hybridized with native Hawaiian ducks (*A. wyvilliana*) and New Zealand grey ducks (*A. superciliosa superciliosa*) (Rhymer and Simberloff, 1996). In addition to these frequently occurring types of impacts, particular invasions can have idiosyncratic but nonetheless devastating effects. For example, the North American cane toad (*Rhinella marina*), brought to Australia in an ill-advised attempt at biological control of introduced beetles on cane, kills threatened native predators such as quolls when they attack the highly toxic toad (Smith and Phillips, 2006).

Invasion biologists, especially recently, have explored ways in which particular invasions affect entire ecosystems (Vitousek and Walker, 1989; Ehrenfeld, 2011; Simberloff, 2011), for instance by changing nutrient cycles, fire or hydrological regimes, or physical structure. Such changes affect many species simultaneously. In Florida, invasion by Australian paperbark (*Melaleuca quinquenervia*) leads to more frequent, hotter lightning-induced fires, to which native plants such as sawgrass (*Cladium jamaicense*) and muhly grass (*Muhlenbergia capillaris*) are maladapted (Serbesoff-King, 2003). This invasion is thus transforming parts of the “river of grass” into patches of paperbark-dominated forest, with consequent changes in the animal community. Introduced plant pathogens that devastate the dominant native vegetation can dramatically modify an entire ecosystem. The Asian chestnut blight fungus (*Cryphonectria parasitica*) spread through eastern North America in the

Domain Editor-in-Chief

Donald R. Zak, University of Michigan

Knowledge Domain

Ecology

Article Type

Commentary

Received: September 2, 2013

Accepted: October 7, 2013

Published: December 4, 2013

early 20th century, removing what had been the dominant tree in many regions, and thereby extinguishing insects host-specific on chestnut (Opler, 1978) as well as changing nutrient cycles (Ellison et al., 2005).

Just as global warming would probably continue for decades even if greenhouse gas emissions were suddenly to be rigorously regulated, so would invasion impacts continue long into the future even if all species introductions were suddenly to cease miraculously today. This is because some introduced species remain quite restricted and innocuous for many years before quite abruptly exploding across the landscape with major impacts (Crooks, 2011), and some impacts take a long time to build to substantial levels. In short, there is an “invasion debt” of impacts that will be experienced in the future of currently harmless introductions that are already present (Essl et al., 2011).

The wealth of invasion impacts already detected, the great increase in research revealing previously undetected impacts, and the growth of trade and travel that bring invasions have induced a pessimism among a segment of the scientific community, who take the attitude that we cannot do much to slow the flood of new invasions or to mitigate the impacts of those that are established, so we should abandon the costly effort and “learn to love them” (e.g., Davis, 2009; Davis et al., 2011; Vince, 2011; Thomas, 2013). They should, according to Marris (2010), be accepted as part of the “new normal.” Among restoration ecologists, a movement has arisen arguing that the combination of biological invasions and global climate change is increasingly making the idea of conserving ecosystems and restoring damaged ones an expensive, hopeless pipe dream and that we should accept the resultant “novel ecosystems” as an inevitable part of a “new ecological world order” (Hobbs, Higgs and Hall, 2013).

The public, although sensitized to particularly dramatic regional invasions such as those of the Burmese python in Florida and the “killer alga” *Caulerpa taxifolia* in the Mediterranean (Meinesz, 1999), does not view invasions in general as a major global change of the order of that caused by anthropogenic climate change or land use change. This relative lack of concern is shown by the lethargic response to the issue by policymakers in most nations, which is reflected in the absence of substantial effective regulations that would stem the flow of invaders. For instance, in the United States (one of the nations in which invasions have received the most attention), one can still legally purchase many of the most ecologically damaging invasive plants at garden centers. Until 2012 one could legally buy, sell, and ship Burmese pythons in the United States. The federal regulation stopping such activity came too late, after the python population had exploded in Florida and gained international public attention. Although damaging invasions are increasingly publicized, pronouncements in scientific journals about a “new normal,” “learn to love them,” and a “new ecological world order,” though minority viewpoints, are avidly publicized by the popular media in much the same way that scientifically credentialed deniers of anthropogenic climate change attract disproportionate attention and justify policymakers in forestalling enacting policies appropriate to the scope and scale of the problem. Momentum is building in many nations to take more aggressive action, but, as with global warming, even the invaders that are already present will generate an increasing range of problems in the future (Essl et al., 2011), at least without a greatly enhanced response.

The contention that we cannot do much to prevent or to mitigate invasions, the chief argument for the “learn to love them” school, is incorrect, but the various effective courses of action are not well known by the public. In fact, there are many ways to stem the arrival of new invaders, to eradicate recent invaders, and to keep well-established ones at low densities, and incremental as well as major advances in relevant technologies have occurred recently. The most efficient approach is to keep invaders out in the first place, and the biggest impediment to doing so is the failure of most nations to restrict planned introductions and to constrict pathways, such as ballast water and untreated wood packing, that inadvertently introduce species. A further problem has been simply maintaining an adequately large corps of trained inspectors of incoming people and cargo. However, New Zealand has achieved notable successes in both of these matters, suggesting that other nations could accomplish a lot if the political will were present (Simberloff, 2013).

The second line of defense is an early detection-rapid response (EDRR) system. Again New Zealand has led the way in developing such a system, an Exotic Pest and Disease Hotline and a staff of trained responders, but few nations have followed. Many invasions have been found early by alert citizens, and this observation has led to mobilization of volunteers to work at the front lines on EDDR systems in both Australia and the United States (Simberloff, 2013).

A variety of methods have successfully eradicated over 1,100 invasions of various taxa to aid conservation (Genovesi, 2011), and many more invasions have been eradicated for agricultural or public health purposes. The complexity of such operations and size of target areas has increased substantially. For instance, goats were eradicated from Santiago Island (58,000 ha) by a combination of the classic Judas goat technique and endocrine injections to prolong estrus (Campbell et al., 2005), while the melon fly (*Bactrocera cucurbitae*) was eradicated from the Ryukyu Archipelago, including Okinawa and Amami, by a combination of the sterile-male and male-annihilation methods (Iwahashi, 1996; Kuba et al., 1996). To date, eradication of aquatic and marine invaders has proven more difficult than eradication of terrestrial ones, although there are recent notable successes, such as eradication of the killer alga *Caulerpa taxifolia* from two regions of California (Anderson, 2005) and the Caribbean black-striped mussel (*Mytilopsis sallei*) from northern Australia (Bax et al., 2001).

If eradication is not attempted or fails, mechanical or physical methods, chemical methods, and biological control can sometimes maintain invasive populations at low levels, and all these technologies have seen recent advances. Mechanical and physical approaches are sometimes impractical because of the high cost of labor, although the use of volunteers or prison work crews has contributed greatly to successful maintenance management in certain cases (Simberloff, 2013). In the wake of *Silent Spring* (Carson, 1962), the use of pesticides and herbicides became a bête noire of conservationists. However, many pesticides and herbicides in use today have few if any non-target impacts and have, in some cases, provided good short- and medium-term control of damaging invaders, including the rescue of the endemic Christmas Island red land crab *Gecarcoidea natalis* from likely extinction by invading yellow crazy ants (*Anoplolepis gracilipes*) (<http://www.deh.gov.au/parks/christmas/fauna/crazy.html>, accessed 8/29/06). However, the expense and evolution of resistance to chemicals work against their effective long-term use, and, in addition, for many chemicals with no known short-term non-target impacts, the effects of chronic exposure are unstudied. Potential non-target impacts have bedeviled biological control as well, although tightened testing procedures for insects introduced to control invasive plants have been greatly improved (Simberloff, 2012). Several insect invasions have also been well controlled by introduced parasitoids or predators, but testing and monitoring procedures for non-target impacts are more lax than those for natural enemies of invasive plants (Simberloff, 2012).

Finally, several novel approaches to eradication and maintenance management of invaders suggest that some currently intractable invasions may become manageable in the future. Striking recent advances include the use of toxic “BioBullets” to kill zebra mussels (Aldridge, Elliott and Moggridge, 2006) and a synthetic mimic of a larval pheromone to attract migrating sea lampreys (Sorensen et al., 2005). In addition, a number of “autocidal” methods are under development in which the genetics or physiology of an invasive species is turned against it, as, for example, the use of transgenes to cause common carp to produce all-male progeny (Bax and Thresher, 2009; Thresher et al., 2013).

In sum, biological invasions will continue to be a major feature of the Anthropocene, with myriad and diverse far-reaching impacts. However, their advance can be slowed and their effects mitigated by political will and ongoing progress in invasion biology and management technologies (Simberloff et al., 2013).

References

- Aldridge DC, Elliott P, Moggridge GD. 2006. Microencapsulated BioBullets for the control of biofouling zebra mussels. *Environ. Sci. Technol.* **40**: 975–979.
- Anderson LWJ. 2005. California’s reaction to *Caulerpa taxifolia*: a model for invasive species rapid response. *Biol. Invasions* **7**: 1003–1016.
- Bax N, Hewitt C, Campbell M, Thresher R. 2001. Man-made marinas as sheltered islands for alien marine organisms: establishment and world-first eradication of a marine pest, in Clout M, Veitch D, eds. *Turning the Tide: Eradication of Invasive Species*. Auckland, NZ: World Conservation Union [IUCN] Species Survival Commission: p. 8.
- Bax NJ, Thresher RE. 2009. Ecological, behavioral and genetic factors influencing the recombinant control of invasive pests. *Ecol. Applic.* **19**: 873–888.
- Beggs JR, Wardle D. 2006. Keystone species: competition for honeydew among exotic and indigenous species, in Allen RB, Lee WG, eds. *Biological Invasions in New Zealand*. Berlin: Springer: p. 281–294.
- Campbell KJ, Baxter GS, Murray PJ, Coblenz BE, Donlan CE, et al. 2005. Increasing the efficacy of Judas goats by sterilization and pregnancy termination. *Wildl. Res.* **32**: 737–743.
- Carson R. 1962. *Silent Spring*. Boston: Houghton Mifflin.
- Cronk QCB. 1989. The past and present vegetation of St Helena. *J. Biogeogr.* **16**: 47–64.
- Crooks JA. 2011. Lag times, in Simberloff D, Rejmánek M, eds. *Encyclopedia of Biological Invasions*. Berkeley, CA: Univ of California Press: p. 404–410.
- Davis MA. 2009. *Invasion Biology*. New York: Oxford Univ Press.
- Davis MA, Chew MK, Hobbs RJ, Lugo AE, Ewel JJ, et al. 2011. Don’t judge species on their origins. *Nature* **474**: 153–154.
- Dorcas ME, Willson JD, Reed RN, Snow RW, Rochford MR, et al. 2012. Severe mammal declines coincide with proliferation of invasive Burmese pythons in Everglades National Park. *Proc. Nat. Acad. Sci (USA)*. **109**: 2418–2422.
- Ehrenfeld JG. 2011. Ecosystem consequences of biological invasions. *Annu. Rev. Ecol. Syst.* **41**: 59–80.
- Ellison AM, Bank MS, Clinton BD, Colburn EA, Elliott K, et al. 2005. Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment* **9**: 479–486.
- Essl F, Dullinger S, Rabitsch W, Hulme PE, Hülber K, et al. 2011. Socioeconomic legacy yields an invasion debt. *Proc. Nat. Acad. Sci (USA)*. **108**: 203–207.
- Genovesi P. 2011. Eradication, in Simberloff D, Rejmánek M, eds. *Encyclopedia of Biological Invasions*. Berkeley: Univ of California Press: p. 198–203.
- Hobbs RJ, Higgs ES, Hall CM, eds. 2013. *Novel Ecosystems: Intervening in the New Ecological World Order*. Oxford, UK: John Wiley & Sons.
- Iwahashi O. 1996. Problems encountered during long-term SIT program in Japan, in McPherson BA, Steck GJ, eds. *Fruit Fly Pests: A World Assessment of Their Biology and Management*. Delray Beach, FL: St. Lucie Press: p. 391–398.
- Kuba H, Kohama T, Kakinohana H, Yamagishi M, Kinjo K, et al. 1996. The successful eradication programs of the melon fly in Okinawa, in McPherson BA, Steck GJ, eds. *Fruit Fly Pests: A World Assessment of Their Biology and Management*. Delray Beach, FL: St. Lucie Press: p. 534–550.
- Lodge DM, Deines A, Gherardi F, Yeo DCJ, Arcella T, et al. 2012. Global introductions of crayfishes: evaluating the impact of species invasions on ecosystem services. *Annual Review of Ecology, Evolution, and Systematics* **43**: 449–472.

- Marris E. 2010. The new normal. *Conservation Magazine* 11: 13–17.
- Meinesz A. 1999. *Killer Algae*. Chicago: University of Chicago Press.
- Opler PA. 1978. Insects of American chestnut: possible importance and conservation concern, in McDonald J, ed. *The American Chestnut Symposium*. Morgantown, WV: West Virginia University Press: 83–85.
- Pascal M. 2011. Rats, in: Simberloff D, Rejmánek M, eds. *Encyclopedia of Biological Invasions*. Berkeley: Univ of California Press: p. 571–575.
- Rhymer J, Simberloff D. 1996. Extinction by hybridization and introgression. *Annu. Rev. Ecol Syst.* 27: 83–109.
- Rushton SP, Lurz PWW, Gurnell J, Nettleton P, Bruemmer C, et al. 2006. Disease threats posed by alien species: the role of a poxvirus in the decline of the native red squirrel in Britain. *Epidemiology and Infection* 134: 521–533.
- Serbesoff-King K. 2003. *Melaleuca* in Florida: a literature review on the taxonomy, distribution, biology, ecology, economic importance and control measures. *J. Aquat. Plant Mgt.* 41: 98–112.
- Simberloff D. 2011. How common are invasion-induced ecosystem impacts? *Biol Invasions* 13: 1255–1268.
- Simberloff D. 2012. Risks of biological control for conservation purposes. *BioControl* 57: 263–276.
- Simberloff D. 2013. Biological invasions: What's worth fighting for and what can be won? *Ecological Engineering*, in press.
- Simberloff D, Martin J-L, Genovesi P, Maris V, Wardle DA, et al. 2013. Impacts of biological invasions: what's what and the way forward. *Trends Ecol. Evol.* 28: 58–66.
- Smith JG, Phillips BL. 2006. Toxic tucker: the potential impact of cane toads on Australian reptiles. *Pacific Conserv Biol* 12: 40–49.
- Sorensen PW, Fine JM, Dvornikovs V, Jeffrey CS, Shao F, et al. 2005. Mixture of new sulfated steroids functions as a migratory pheromone in the sea lamprey. *Nature Chemical Biology* 1: 324–328.
- Thomas CD. 2013. The Anthropocene could raise biological diversity. *Nature* 502: 7.
- Thresher R, Hayes K, Bax N, Teem J, Benfey T, et al. 2013. Genetic control of invasive fish: technological options and its role in Integrated Pest Management. *Biol. Invasions*, in press.
- Turner GG, Reeder DM, Coleman JTH. 2011. A five-year assessment of mortality and geographic spread of white-nose syndrome in North American bats, with a look at the future. Update of white-nose syndrome in bats. *Bat Research News* 52: 13–27.
- van Riper CIII, van Riper SG, Goff ML, Laird M. 1986. The epizootiology and ecological significance of malaria in Hawaiian land birds. *Ecol Monogr* 56: 327–344.
- Vince G. 2011. Embracing invasives. *Science* 331: 1383–1384.
- Vitousek PM, Walker LR. 1989. Biological invasion by *Myrica faya* in Hawai'i: plant demography, nitrogen fixation, ecosystem effects. *Ecol. Monogr* 59: 247–265.
- Wauters LA, Tosi G, Gurnell J. 2002. Interspecific competition in tree squirrels: do introduced grey squirrels (*Sciurus carolinensis*) deplete tree seeds hoarded by red squirrels (*S. vulgaris*)? *Behav Ecol Sociobiol* 51: 360–367.

Copyright

© 2013 Simberloff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.