

REVIEW

Pharmaceuticals and personal care products (PPCPs) are ecological disrupting compounds (EcoDC)

Erinn K. Richmond*, Michael R. Grace*, John J. Kelly[†], Alexander J. Reisinger[‡], Emma J. Rosi[§] and David M. Walters^{||}

Pharmaceuticals and personal care products (PPCPs) are ubiquitous in freshwater ecosystems worldwide and are recognized as contaminants of concern. Currently, contaminants of concern are classified for their persistence, bioaccumulation, and toxicity (PBT criteria). PPCPs are not classified as persistent organic pollutants (POPs), although some PPCPs share characteristics similar to POPs. For example, PPCPs are known to be pseudopersistent due to constant discharge into the environment, often at low concentrations. At commonly reported environmental concentrations, PPCPs are rarely toxic, but the ability of these compounds to disrupt ecological processes and functions in freshwater ecosystems is often overlooked. Herein we briefly summarize recent studies highlighting the potential ecological effects of PPCPs, including effects on key ecological processes (e.g. primary productivity and community respiration), and we propose that appropriate screening for harmful effects of PPCPs in surface waters should be expanded to include Ecologically Disrupting Compounds (EcoDC) in addition to the established PBT criteria.

Keywords: PPCPs; ecological processes; disruption; pharmaceuticals; sub-lethal; EcoDC

Introduction

Pharmaceuticals and personal care products (PPCPs) are biologically active compounds that are recognized as environmental contaminants of global concern due to their presence in ecosystems throughout the world (Monteiro and Boxall, 2010) and more recently as agents of global change (Bernhardt et al., 2017). PPCPs can enter the environment through multiple pathways, including both point and non-point sources, and are commonly detected in surface waters at low concentrations (ng to $\mu\text{g L}^{-1}$) (Daughton and Ternes, 1999). As the global human population and percentage of the population living in high-density urban areas continue to increase, PPCP contamination of ecosystems is expected to increase substantially both in the number of contaminated ecosystems, and in the typical PPCP concentration found in the environment (Weber et al., 2015). Additionally, it is likely that the contribution of increased water usage to global water shortages will

further exacerbate concentrations of PPCPs (Klaminder et al., 2015; Petrovic et al., 2011).

Relative to other types of organic chemicals, e.g. polychlorinated biphenyls (PCBs), questions surrounding the biological impacts of PPCPs remain largely unanswered (Monteiro and Boxall, 2010). Currently, organic contaminants are screened and classified for potential harmful effects using PBT (persistence, bioaccumulation, and toxicity) criteria (EPA, 2012; Klečka et al., 2009; Stempel et al., 2012). In contrast to the US EPA approach, the European Union Water Framework Directive has identified a list of priority substances based upon similar criteria to the EPA, but using precautionary principles, this list is to be updated every 4 years to include pseudopersistent PPCPs (Ellis, 2006). Although many PPCPs have a limited lifespan in the environment, PPCPs are often classified as pseudopersistent based on the continual addition of these compounds to the environment (Daughton and Ternes, 1999). Furthermore, our knowledge of the persistence of PPCPs in the environment is often based on laboratory studies, which can underestimate persistence for certain compounds (Klaminder et al., 2015). A review by Daughton and Ternes (1999) highlighted the need to study the environmental fate of PPCPs in a similar fashion to which persistent organic pollutants (POPs) such as some pesticides and other organic contaminants have been studied. Despite this call for more research on the environmental fate of PPCPs, we still lack a solid understanding of the fate and bioaccumulation of PPCPs in the environment (e.g. Walters et al., 2016) and of the effects

* Water Studies Centre, School of Chemistry, Monash University, Clayton Victoria, AU

[†] Department of Biology, Loyola University, Chicago, Illinois, US

[‡] Soil and Water Sciences Department, University of Florida, Gainesville, Florida, US

[§] Cary Institute of Ecosystem Studies, Millbrook, New York, US

^{||} U.S. Geological Survey, Fort Collins, Colorado, US

Corresponding author: Erinn K. Richmond
(erinn.richmond@monash.edu)

of PPCPs on ecosystem function. This knowledge gap is particularly noteworthy, as synthetic chemicals similar to PPCPs have recently received attention and are also not well studied in ecological and environmental research (Bernhardt et al., 2017).

In 1999 the United States Geological Survey (USGS) undertook a national survey to quantify the presence of organic contaminants, including PPCPs, in US surface waters. The results of this study indicated that organic contaminants were detected in 80% of streams sampled across 30 US states and that detection of multiple contaminants in a single sample was common (Kolpin et al., 2002). Similar results were observed in a recent survey conducted by the USGS, where 84 PPCPs were detected in 38 US streams (Bradley et al., 2017). The landmark study by Kolpin et al. (2002), emulated by numerous additional studies, highlighted the many PPCPs in surface waters throughout the world (e.g. Stewart et al., 2014; Kasprzyk-Hordén et al., 2008). A majority of these compounds, largely understudied, are currently unregulated or have been determined to be low risk due to low environmental concentrations, possibly because the perceived social benefit outweighs associated environmental risks. Much of the research and regulatory requirements exploring the effects of these compounds have used traditional toxicity tests (e.g. the concentration required for PPCPs to cause 50% mortality (LC50s) in model organisms, or the predicted no effect concentration (PNEC) of a compound). These conventional toxicity tests included in the PBT criteria show that PPCPs have the capacity to be toxic at high concentrations (exceeding human therapeutic doses) and typically conclude that PPCPs in the environment are not hazardous due to low environmental concentrations, despite often only testing effects on model organisms (e.g. the common green alga *Scenedesmus*). More simply, the LC50s of these compounds are much higher than concentrations detected in the environment. Therefore, pharmaceuticals are usually classified as non-toxic, although a notable publicized exception was the widespread mortality of old world vultures consuming dead livestock that had been previously treated with a veterinary anti-inflammatory drug (Oaks et al., 2004).

Despite the general lack of toxic effects at environmentally relevant concentrations, PPCPs are capable of causing various sub-lethal ecological effects on many components of aquatic ecosystems, thus emphasis should be placed on evaluating over all ecosystem effects of PPCPs. The US Food and Drug Administration (FDA) only requires an environmental assessment report for new drug applications if the concentration reaching surface waters is $>1 \mu\text{g/L}$ (FDA, 1998). However, pharmaceuticals are designed to be effective and used at low therapeutic doses, potentially causing sub-lethal effects in natural environments. Therefore, we contend that the effects of these compounds cannot be readily ascertained with traditional toxicity testing alone and that the current emphasis on single organismal lethality could lead the scientific and regulatory community to underestimate the potential risks of PPCPs to aquatic ecosystems.

Known ecological disrupting effects of PPCPs

Recent research investigating sub-lethal ecological effects of PPCPs has demonstrated that environmentally relevant concentrations of these compounds may alter ecological interactions and processes (**Table 1**). PPCPs can also alter relationships among organisms. For example, a common antidepressant, oxazepam, alters the feeding behavior of European perch (Brodin et al., 2013; Brodin et al., 2014), and tadpoles (*Bufo arabicus*) were more susceptible to predation from predatory dragonfly larvae (*Anax imperator*) when exposed to low concentrations of the selective serotonin reuptake inhibitor (SSRI) fluoxetine ($3 \mu\text{g/L}$) (Barry, 2014). PPCPs can also promote changes in ecosystem structure and function, including biofilm primary production, respiration and community structure, biogeochemical processes, and invertebrate growth and population dynamics. Microbial respiration was suppressed when exposed to common pharmaceuticals including caffeine (53%), ciprofloxacin (91%), cimetidine (51%) and diphenhydramine (63%) which also almost completely suppressed photosynthesis (99%) (Rosi-Marshall et al., 2013). Similarly, biofilm primary production was suppressed by over 88% when exposed to the illicit drug amphetamine (Lee et al., 2016). In addition, fluoxetine and citalopram, which are commonly used antidepressants, reduced algal production by 29% (Richmond et al., 2016). Altering one or several ecosystem processes can irreparably disrupt ecological systems. For example decreases in primary production can reduce the organic matter available to higher trophic levels, thereby altering food web dynamics (Polis and Strong, 1996).

Similar to functional responses, PPCPs can induce alterations to bacterial and diatom communities and can select for antimicrobial resistance both in laboratory and field settings. Diphenhydramine exposure altered bacterial composition on novel test substrates in streams by increasing the relative abundance of a common biofilm producing bacterium, *Pseudomonas*, while causing relative decreases in *Flavobacterium* sp. (Rosi-Marshall et al., 2013). Likewise, artificial streams amended with $1 \mu\text{g/L}$ amphetamine significantly altered diatom communities (Lee et al., 2016), and a combination of 9 PPCPs (total $5 \mu\text{g/L}$) and altered flow regimes reduced biofilm taxonomic richness and biomass, while also changing the structure of the bacterial community (Corcoll et al., 2015). Furthermore, concentrations of triclosan, a common antimicrobial, have been found to alter the structure of bacterial communities and increase antimicrobial resistant bacteria (Drury et al., 2013b; Proia et al., 2011), both in situ and in artificial streams amended with triclosan. In addition, streams that receive wastewater effluent displayed significantly reduced abundance and diversity of benthic bacterial communities (Drury et al., 2013a), although this may also have been as a result of other contaminants found in wastewater besides PPCPs.

Pharmaceuticals can also disrupt and alter processes within aquatic invertebrate communities, and these disruptions or alterations can in turn disrupt other ecosystem functions. For example, aquatic insect metamorphosis and emergence occurred sooner and at

Table 1: Examples of relevant studies documenting ecological disruption resulting from low-dose, sub-lethal environmental concentrations of PPCPs. DOI: <https://doi.org/10.1525/elementa.252.t1>

PPCP	Test biota	Concentration ($\mu\text{g/L}^{-1}$)	Effect summary	Citation
Amphetamine (illicit drug)	Stream benthic communities	1	Suppression of gross primary production on autotrophic biofilms, compositional shift of bacterial and biofilm communities, increased dipteran (stream insect) emergence	(Lee et al., 2016)
Caffeine (stimulant), acetaminophen (pain reliever), diclofenac (anti-inflammatory)	Protozoa, micrometazoa and algae	5	Increased abundance of organisms and feeding activity (direct), decreased biofilm (algal) biomass (indirect)	(Lawrence et al., 2012)
Cimetidine (antihistamine)	Invertebrates (<i>Gammarus fasciatus</i> and <i>Psephenus herricki</i>)	0.07–70	Reduced growth and biomass of <i>G. fasciatus</i> , low survivorship of <i>P. herricki</i> when exposed to high concentrations	(Hoppe et al., 2012)
Diuron (herbicide), triclosan (antimicrobial)	River biofilm communities	15–60	Biofilm recovery after exposure to short term pulses of triclosan and diuron	(Proia et al., 2011)
Fluoxetine (antidepressant), citalopram (antidepressant)	Algal and invertebrate benthic communities	20	Suppression of primary productivity and community respiration on biofilms (algae). Increased stream insect emergence	(Richmond et al., 2016)
Oxazepam (antidepressant)	European Perch (<i>Perca fluviatilis</i>)	1.8	Increased feeding rates and locomotor activity (direct); implications for prey populations (indirect)	(Brodin et al., 2013)
Oxazepam (antidepressant)	European Perch (<i>Perca fluviatilis</i>), Mayfly (<i>Coenagrion hastulatum</i>)	2	Increased predator activity of perch, no behavioral effects on mayflies (direct). Increased bioaccumulation via consumption of exposed prey (indirect)	(Brodin et al., 2014)
Triclosan (anti-microbial)	Benthic bacterial communities	1–107 (field sites) 17.3 $\mu\text{g/g}$ (max concentration, mesocosm)	Field and laboratory experiment. Correlation between sediment triclosan concentration and abundance of triclosan resistant bacteria at field sites. Increased abundance of triclosan resistant bacteria in mesocosms	(Drury et al., 2013b)
Triclosan (antimicrobial)	Stream benthic communities	0.1–10	Increase in abundance of triclosan resistant bacteria and stimulation of periphyton growth	(Nietch et al., 2013)
Wastewater treatment plant (WWTP) effluent	Benthic bacterial communities	N/A	Field experiment. Changes in bacterial abundance and community composition above and below WWTPs	(Drury et al., 2013a)

a greater rate when insect larvae (Diptera) were exposed to 20 µg/L of fluoxetine and citalopram (Richmond et al., 2016), and cumulative insect emergence increased by up to 89% when exposed to amphetamine (Lee et al., 2016). Cimetidine, a common antihistamine, reduced *Gammarus* biomass when exposed to low concentrations (0.07–70 µg/L) (Hoppe et al., 2012) and exposure to low concentrations of an antibiotic mixture (2 µg/L) led to shifts in leaf microbial communities, resulting in increased amphipod (*Gammarus*) body mass (Bundschuh et al., 2017). Changes in animal biomass, particularly for primary consumers near the base of a food web, can affect population and community dynamics within aquatic ecosystems, whereas altered phenology and mass fluxes of aquatic insect emergence have energetic implications for linked aquatic and terrestrial food webs (Baxter et al., 2005), and can alter riparian community structure (Kalcounis-Rueppell et al., 2007).

It is also worth noting that ecological disruption by PPCPs can be direct or indirect, analogous to other traditional drivers of ecological interactions (McQueen et al., 1986; Polis et al., 1997; Wootton, 1994). For example, as mentioned earlier, European Perch (*Perca fluviatilis*) exposed to low concentrations (1.8 µg/L) of oxazepam, a common psychoactive drug, accumulated the pharmaceutical in their tissues and exhibited altered behavior, including increased feeding rate (direct effect; Brodin et al., 2013). An increase in perch feeding rate could affect prey populations (indirect effect), even if the prey species was not directly affected by the drug exposure (Brodin et al., 2014). Similarly, caffeine stimulates the numbers and grazing activity of microscopic animals such as nematodes and rotifers (direct effect), which results in decreased biomass of algae and bacteria in aquatic biofilms (indirect effect; Lawrence et al., 2012). These indirect ecological effects demonstrate that a specific biochemical interaction is not always necessary for a pharmaceutical to act as an ecological disruptor and affect organisms, community dynamics, or ecological processes (Table 1). Moreover, the ecological effects of single PPCP compound exposure may be incalculably increased by simultaneous exposure to the wide range of PPCP compounds commonly detected in surface waters (Bradley et al., 2017). Many of these effects remain largely unknown or undetected, in part because they may be subtle and difficult to measure. Based upon this recent research, we conclude that PPCPs can have profound ecological impacts capable of disrupting ecological dynamics. Therefore, we propose that appropriate screening for harmful effects of PPCPs in surface waters should be expanded to include ecological disruption, highlighting PPCPs as Ecological Disrupting Compounds (EcoDC), in addition to the established PBT criteria.

Our concept of EcoDCs (Figure 1A) is analogous to endocrine disrupting compounds (EDCs), which are generally non-toxic but can alter the endocrinology of organisms at environmentally relevant concentrations (Clotfelter et al., 2004). The presence of EDCs in surface waters has been well documented over the last two decades, and a growing body of literature has examined the potential for EDCs to cause non-lethal effects on aquatic biota at

low concentrations, particularly changes in behavior and sexual reproduction of organisms (Clotfelter et al., 2004; Colborn et al., 1993). In 1998, the EPA Endocrine Disruptor Screening and Testing Advisory Committee was formed to make federal recommendations regarding adverse effects of EDCs found in the environment (EPA, 1998). After the recommendations of the committee were published, the presence of EDCs in the environment and their ability to cause non-lethal effects has received considerable attention.

Research priorities and implications for resource and policy managers

Despite the need for a better understanding of the environmental fate of PPCPs (Daughton and Ternes, 1999) and a growing body of evidence documenting sub-lethal and ecosystem-level effects, studies on toxicity of PPCPs vastly outnumber those on ecosystem disruption. Based on a literature search using Web of Science, >90% of scientific articles studying the effects of PPCPs have quantified lethal endpoints (Figure 1B). Of the articles that measured endpoints associated with ecological disruption at environmentally relevant PPCP concentrations, over half (55%) focused on behavioral alterations of a target organism, rather than quantifying changes in ecological functions and community dynamics or indirect effects (Figure 1C). The apparent lack of contaminant studies focusing on ecological endpoints can be at least partially attributed to the limited funding for research on PPCPs by major funding agents such as the National Science Foundation (Bernhardt et al., 2017). This funding gap, coupled with the focus on traditional toxicity testing, results in a loss of cohesion between ecotoxicological and ecological disciplines and thus long term, subtle, ecological disruption endpoints are often overlooked.

The use of PPCPs will likely continue to increase throughout the world as the proportion of the global population with access to treatment increases, so the release of PPCPs to surface waters is likely to increase rather than abate in the near future (Schroder et al., 2016). In addition, increased water-use efficiency will likely contribute to an increase of PPCP concentrations in the environment (Klaminder et al., 2015). Although the toxic effects of many PPCPs are limited to concentrations much higher than typically detected in the environment, there is growing scientific evidence that these compounds can act as Ecological Disrupting Compounds with heretofore unknown and unpredictable long-term effects. Current chemical and contaminant management practices do not appropriately address these issues and we argue that an additional standardized criterion should be established for risk assessments of these biologically active compounds that includes consideration of their potential to disrupt ecological processes across multiple components of an aquatic ecosystem, and that this criterion recognizes this potential even at low (<1 µg/L) concentrations. Beyond the scope of the current review, efforts to develop standardized protocols to quantify sub-lethal effects as well as community- and ecosystem-level effects are needed. Although difficult to standardize due to the myriad




Figure 1: PPCPs as EcoDCs influence aquatic ecosystem function and biota. A) Conceptual figure demonstrating the potential non-lethal effects of EcoDCs at low concentrations on aquatic biota and ecosystem function, in contrast to toxic effects observed at high concentrations. The dashed horizontal line at the end of this curve represents uncertainty at high PPCP concentrations. The dashed vertical line indicated where lethality is becoming increasingly important relative to ecosystem disruption. **B)** Number of publications indexed in Web of Science measuring toxic endpoints (including LC50 and EC50 toxicity testing) versus those measuring non-lethal, ecological disruption (EcoD) endpoints at environmentally relevant concentrations. **C)** Number of publications in Web of Science demonstrating various endpoints related to ecosystem disruption associated with environmentally relevant concentrations of PPCPs. Details of literature search are available in supplementary material. DOI: <https://doi.org/10.1525/elementa.252.f1>

environmental conditions that could affect PPCP EcoDC, future research and protocol development in this area are vital. Building off of this argument, we recommend adding EcoDC to the PBT toxicology criteria for this group of compounds. In support of this recommendation we have provided examples to demonstrate that pharmaceuticals have biological consequences at low, non-toxic concentrations. These concentrations seldom cause mortality, but

have the ability to alter life histories and behaviors, and change processes and community structure within aquatic systems. We further suggest that future research consider testing for ecological disruption and focus on the potential effects of pharmaceuticals on the ecology and biology at all levels of aquatic ecosystems to better understand the long-term consequences of environmental exposures to these organic chemicals.

Supplemental File

The supplemental file for this article can be found as follows:

- **Text S1.** Search terms in Web of Science for Figure 1B and C. DOI: <https://doi.org/10.1525/elementa.252.s1>

Acknowledgements

We thank Dr. Emily Bernhardt for providing helpful comments on an earlier draft of the manuscript. This research was subjected to USGS review and approved for publication. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Funding information

We acknowledge that funding for Erinn Richmond was provided by Monash University, Faculty of Science Dean's Postgraduate award.

Competing interests

The authors have no competing interests to declare.

Author contributions

- Contributed to conception and design: EKR, MRG, JJK, AJR, EJR, DMW
- Contributed to acquisition of data: EKR, EJR, DMW
- Contributed to analysis and interpretation of data: EKR, EJR, DMW
- Drafted and/or revised the article: EKR, MRG, JJK, AJR, EJR, DMW
- Approved the submitted version for publication: EKR, MRG, JJK, AJR, EJR, DMW

References

- Barry, MJ** 2014 Fluoxetine inhibits predator avoidance behavior in tadpoles. *Toxicological & Environmental Chemistry* **96**(4): 641–649. DOI: <https://doi.org/10.1080/02772248.2014.966713>
- Baxter, CV, Fausch, KD and Saunders, CW** 2005 Tangled webs: reciprocal flows of invertebrate prey link streams and riparian zones. *Freshwater Biology* **50**(2): 201–220. DOI: <https://doi.org/10.1111/j.1365-2427.2004.01328.x>
- Bernhardt, ES, Rosi, EJ and Gessner, MO** 2017 Synthetic chemicals as agents of global change. *Frontiers in Ecology and the Environment*, 84–90. DOI: <https://doi.org/10.1002/fee.1450>
- Bradley, PM, Journey, CA, Romanok, KM, Barber, LB, Buxton, HT, et al.** 2017 Expanded target-chemical analysis reveals extensive mixed-organic-contaminant exposure in U.S. streams. *Environmental Science & Technology* **51**(9): 4792–4802. DOI: <https://doi.org/10.1021/acs.est.7b00012>
- Brodin, T, Fick, J, Jonsson, M and Klaminder, J** 2013 Dilute concentrations of a psychiatric drug alter behavior of fish from natural populations. *Science* **339**(6121): 814–815. DOI: <https://doi.org/10.1126/science.1226850>
- Brodin, T, Piovano, S, Fick, J, Klaminder, J, Heynen, M, et al.** 2014 Ecological effects of pharmaceuticals in aquatic systems—impacts through behavioural alterations. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**(1656). DOI: <https://doi.org/10.1098/rstb.2013.0580>
- Bundschuh, M, Hahn, T, Gessner, MO and Schulz, R** 2017 Antibiotic mixture effects on growth of the leaf-shredding stream detritivore *Gammarus fossarum*. *Ecotoxicology* **26**(4): 547–554. DOI: <https://doi.org/10.1007/s10646-017-1787-2>
- Clotfelter, ED, Bell, AM and Levering, KR** 2004 The role of animal behaviour in the study of endocrine-disrupting chemicals. *Animal Behaviour* **68**(4): 665–676. DOI: <https://doi.org/10.1016/j.anbehav.2004.05.004>
- Colborn, T, vom Saal, FS and Soto, AM** 1993 Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environmental Health Perspectives* **101**(5): 378–384. DOI: <https://doi.org/10.1289/ehp.93101378>
- Corcoll, N, Casellas, M, Huerta, B, Guasch, H, Acuña, V, et al.** 2015 Effects of flow intermittency and pharmaceutical exposure on the structure and metabolism of stream biofilms. *Science of The Total Environment* **503–504**: 159–170. DOI: <https://doi.org/10.1016/j.scitotenv.2014.06.093>
- Daughton, CG and Ternes, TA** 1999 Pharmaceuticals and personal care products in the environment: agents of subtle change? *Environmental Health Perspectives* **107**: 907–938. DOI: <https://doi.org/10.1289/ehp.99107s6907>
- Drury, B, Rosi-Marshall, E and Kelly, JJ** 2013a Wastewater treatment effluent reduces the abundance and diversity of benthic bacterial communities in urban and suburban rivers. *Applied and Environmental Microbiology* **79**(6): 1897–1905. DOI: <https://doi.org/10.1128/AEM.03527-12>
- Drury, B, Scott, J, Rosi-Marshall, EJ and Kelly, JJ** 2013b Triclosan exposure increases triclosan resistance and influences taxonomic composition of benthic bacterial communities. *Environmental Science & Technology* **47**(15): 8923–8930. DOI: <https://doi.org/10.1021/es401919k>
- Ellis, JB** 2006 Pharmaceutical and personal care products (PPCPs) in urban receiving waters. *Environmental Pollution* **144**(1): 184–189. DOI: <https://doi.org/10.1016/j.envpol.2005.12.018>
- EPA** 1998 Endocrine disruptor screening and testing advisory committee final report – executive summary. <https://www.epa.gov/sites/production/files/2015-08/documents/exesum14.pdf> Accessed 8/4/16.
- EPA** 2012 Persistent, bioaccumulative, and toxic profiles estimated for organic chemicals on-line. <http://www.pbtprofiler.net/> Accessed 8/4/16.
- FDA** 1998 Guidance for industry – Environmental impact assesment of human drug and biologica

- applications, Revision 1. Rockville. <https://www.fda.gov/downloads/Drugs/Guidances/ucm070561.pdf> Accessed 8/4/16.
- Hoppe, PD, Rosi-Marshall, EJ and Bechtold, HA** 2012 The antihistamine cimetidine alters invertebrate growth and population dynamics in artificial streams. *Freshwater Science* **31**(2): 379–388. DOI: <https://doi.org/10.1899/11-089>
- Kalcounis-Rueppell, MC, Payne, VH, Huff, SR and Boyko, AL** 2007 Effects of wastewater treatment plant effluent on bat foraging ecology in an urban stream system. *Biological Conservation* **138**(1–2): 120–130. DOI: <https://doi.org/10.1016/j.biocon.2007.04.009>
- Kasprzyk-Hordern, B, Dinsdale, RM and Guwy, AJ** 2008 The occurrence of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs in surface water in South Wales, UK. *Water Research* **42**(13): 3498–3518. DOI: <https://doi.org/10.1016/j.watres.2008.04.026>
- Klaminder, J, Brodin, T, Sundelin, A, Anderson, NJ, Fahlman, J, et al.** 2015 Long-term persistence of an anxiolytic drug (oxazepam) in a large freshwater lake. *Environmental Science & Technology* **49**(17): 10406–10412. DOI: <https://doi.org/10.1021/acs.est.5b01968>
- Klečka, GM, Muir, DCG, Dohmen, P, Eisenreich, SJ, Gobas, FAPC, et al.** 2009 Introduction to special series: Science-based guidance and framework for the evaluation and identification of PBTs and POPs. *Integrated Environmental Assessment and Management* **5**(4): 535–538. DOI: https://doi.org/10.1897/IEAM_2009-045.1
- Kolpin, DW, Furlong, ET, Meyer, MT, Thurman, EM, Zaugg, SD, et al.** 2002 Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams, 1999–2000: A national reconnaissance. *Environmental Science & Technology* **36**(6): 1202–1211. DOI: <https://doi.org/10.1021/es011055j>
- Lawrence, JR, Zhu, B, Swerhone, GDW, Roy, J, Tumber, V, et al.** 2012 Molecular and microscopic assessment of the effects of caffeine, acetaminophen, diclofenac, and their mixtures on river biofilm communities. *Environmental Toxicology and Chemistry* **31**(3): 508–517. DOI: <https://doi.org/10.1002/etc.1723>
- Lee, SS, Paspalof, AM, Snow, D, Richmond, EK, Rosi-Marshall, EJ, et al.** 2016 Occurrence and potential biological effects of amphetamine on stream communities. *Environmental Science & Technology* **50**(17): 9727–35. DOI: <https://doi.org/10.1021/acs.est.6b03717>
- McQueen, DJ, Post, JR and Mills, EL** 1986 Trophic Relationships in Freshwater Pelagic Ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences* **43**(8): 1571–1581. DOI: <https://doi.org/10.1139/f86-195>
- Monteiro, S and Boxall, AA** 2010 Occurrence and Fate of Human Pharmaceuticals in the Environment. *Reviews of Environmental Contamination and Toxicology*, 53–154. Springer: New York.
- Nietch, CT, Quinlan, EL, Lazorchak, JM, Impellitteri, CA, Raikow, D, et al.** 2013 Effects of a chronic lower range of triclosan exposure on a stream mesocosm community. *Environmental Toxicology and Chemistry* **32**(12): 2874–2887. DOI: <https://doi.org/10.1002/etc.2385>
- Oaks, JL, Gilbert, M, Virani, MZ, Watson, RT, Meteyer, CU, et al.** 2004 Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature* **427**(6975): 630–633. http://www.nature.com/nature/journal/v427/n6975/supinfo/nature02317_S1.html. DOI: <https://doi.org/10.1038/nature02317>
- Petrovic, M, Ginebreda, A, Acuña, V, Batalla, RJ, Elosegi, A, et al.** 2011 Combined scenarios of chemical and ecological quality under water scarcity in Mediterranean rivers. *TrAC Trends in Analytical Chemistry* **30**(8): 1269–1278. DOI: <https://doi.org/10.1016/j.trac.2011.04.012>
- Polis, GA, Anderson, WB and Holt, RD** 1997 Toward an integration of landscape and food web ecology: the dynamics of spatially subsidized food webs. *Annual Review of Ecology and Systematics* **28**: 289–316. DOI: <https://doi.org/10.1146/annurev.ecolsys.28.1.289>
- Polis, GA and Strong, DR** 1996 Food web complexity and community dynamics. *The American Naturalist* **147**(5): 813–846. DOI: <https://doi.org/10.1086/285880>
- Proia, L, Morin, S, Peipoch, M, Romaní, AM and Sabater, S** 2011 Resistance and recovery of river biofilms receiving short pulses of Triclosan and Diuron. *Science of The Total Environment* **409**(17): 3129–3137. DOI: <https://doi.org/10.1016/j.scitotenv.2011.05.013>
- Richmond, EK, Rosi-Marshall, EJ, Lee, SS, Grace, MR and Thompson, RM** 2016 Antidepressants in stream ecosystems: Influence of selective serotonin reuptake inhibitors (SSRIs) on algal production and insect emergence. *Freshwater Science* **35**(3): 845–855. DOI: <https://doi.org/10.1086/687841>
- Rosi-Marshall, EJ, Kincaid, DW, Bechtold, HA, Royer, TV, Rojas, M, et al.** 2013 Pharmaceuticals suppress algal growth and microbial respiration and alter bacterial communities in stream biofilms. *Ecological Applications* **23**(3): 583–593. DOI: <https://doi.org/10.1890/12-0491.1>
- Schröder, P, Helmreich, B, Škrbić, B, Carballa, M, Papa, M, et al.** 2016 Status of hormones and painkillers in wastewater effluents across several European states—considerations for the EU watch list concerning estradiols and diclofenac. *Environmental Science and Pollution Research* **23**(13): 12835–12866. DOI: <https://doi.org/10.1007/s11356-016-6503-x>
- Stewart, M, Olsen, G, Hickey, CW, Ferreira, B, Jelić, A, et al.** 2014 A survey of emerging contaminants

in the estuarine receiving environment around Auckland, New Zealand. *Science of The Total Environment* **468–469**: 202–210. DOI: <https://doi.org/10.1016/j.scitotenv.2013.08.039>

Strempel, S, Scheringer, M, Ng, CA and Hungerbühler, K 2012 Screening for PBT Chemicals among the “Existing” and “New” Chemicals of the EU. *Environmental Science & Technology* **46**(11): 5680–5687. DOI: <https://doi.org/10.1021/es3002713>

Walters, DM, Jardine, TD, Cade, BS, Kidd, KA, Muir, DCG, et al. 2016 Trophic magnification of organic

chemicals: A global synthesis. *Environmental Science & Technology* **50**(9): 4650–4658. DOI: <https://doi.org/10.1021/acs.est.6b00201>

Weber, FA, Bergmann, A, Hickmann, S, Ebert, I, Hein, A, et al. 2015 Pharmaceuticals in the environment—global occurrences and perspectives. *Environmental Toxicology and Chemistry*.

Wootton, JT 1994 The nature and consequences of indirect effects in ecological communities. *Annual Review of Ecology and Systematics* **25**: 443–466. DOI: <https://doi.org/10.1146/annurev.es.25.110194.002303>

How to cite this article: Richmond, EK, Grace, MR, Kelly, JJ, Reisinger, AJ, Rosi, EJ and Walters, DM 2017 Pharmaceuticals and personal care products (PPCPs) are ecological disrupting compounds (EcoDC). *Elem Sci Anth*, 5: 66. DOI: <https://doi.org/10.1525/elementa.252>

Domain Editor-in-Chief: Donald R. Zak, University of Michigan, US

Associate Editor: Julian D. Olden, University of Washington, US

Knowledge Domain: Ecology

Submitted: 18 May 2017 **Accepted:** 25 September 2017 **Published:** 13 November 2017

Copyright: © 2017 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

